

A PHASE III, PROSPECTIVE, DOUBLE-BLIND, RANDOMIZED, PLACEBO-CONTROLLED TRIAL OF THROMBOLYSIS IN IMAGING-ELIGIBLE, LATE-WINDOW PATIENTS TO ASSESS THE EFFICACY AND SAFETY OF TENECTEPLASE (TIMELESS)

## Enrollment and Randomization Checklist-HUP

<u>Screer</u>	ning (pre-randomization)		
	Informed consent Review inclusion/exclusion criteria and document on CRF Local labs  CMP Heme/CBC (platelets need to be drawn prior to randomization, but not resulted unless there is a suspicion for underlying coagulopathy!)		
	☐ Coags (only if patient is taking an anticoagulant)		
	☐ Urine pregnancy test if of child bearing potential		
	Obtain subject weight: kg		
	Baseline (pre-stroke) mRS:		
	NIHSS:		
	☐ Ischemic Core Volume: mL		
	Mismatch ratio:		
	☐ Mismatch volume: mL		
Rando	<u>mization</u>		
_	<ul> <li>□ Call inpatient pharmacy and ask to speak to the IV pharmacist to alert them so they can begin to prepare: 215-662-2907</li> <li>□ Log into Endpoint: <a href="https://secure.endpointclinical.com/">https://secure.endpointclinical.com/</a></li> <li>○ Follow these steps in order to randomize:</li> <li>■ Click on IWR drag down under Genentech logo, and select screening</li> <li>■ After screening information is complete, go back to IWR drag down and select randomization to</li> </ul>		
	complete randomization process		
	MAKE SURE THE APPROPRIATE SITE IS SELECTED!		
	<ul> <li>Print randomization with kit assignment and fax to pharmacy: 215-349-5841</li> </ul>		
	☐ Place order in EPIC for study drug- please refer to separate document with instructions		
	Obtain study drug and administer to patient		
г		ı —	
	KEY TIME POINTS TO KNOW!	E	Emergency contacts:
	<ol> <li>Randomization MUST occur within 90 minutes of qualifying brain imaging. If the elapsed time between</li> </ol>	E	Endpoint support: 877-810-4786

- Randomization MUST occur within 90 minutes of qualifying brain imaging. If the elapsed time between imaging and randomization exceeds 90 minutes, repeat imaging must be obtained before proceeding with study enrollment.
- 2. STUDY DRUG MUST BE ADMINSTERED PRIOR TO GROIN PUNCTURE!
- **3.** Following study drug administration, neurologic assessments and vital signs should mirror post-tPA administration (Every 15 minutes for first 2 hours, then every 30 minutes for 6 hours, then hourly until 24 hours post administration)

Hotline: 1-888-316-3335

RAPID: 650-388-9767, ext 2

HUP site number: 317790 PAH site number: 319158 PPMC site number: 319156